## NATIONAL STD/AIDS CONTROL PROGRAMME

## QUARTERLY RETURN OF SEXUALLY TRANSMITTED DISEASE INCLUDING HIV/AIDS FOR EPIDEMIOLOGICAL BULLETIN OF THE EPIDEMIOLOGY UNIT OF MINISTRY OF HEALTH

## New episodes of STD/HIV/AIDS reported or treated at STD clinics in Sri Lanka Year 2016 4th Quarter

Disease	New cases or new disease episodes during the quarter			Total new cases or new episodes for calender year upto end of the quarter#		
	Male	Female	Total	Male	Female	Total
HIV Positives <sup>1</sup>	44	14	58	189	62	251
AIDS	6	2	8	42	11	53
Early Syphilis <sup>2</sup>	15	15	30	88	44	132
Late Syphilis <sup>3</sup>	142	75	217	505	288	793
Congenital Syphilis <sup>4</sup>	1	1	2	4	5	9
Gonorrhoea	62	15	77	234	66	300
Ophthalamia Neonatorum <sup>6</sup>	1	0	1	3	1	4
Non specific cervicitis/urethritis	153	451	604	588	1595	2183
Chlamydial infection	4	0	4	8	9	17
Genetal Herpes	299	418	717	1299	1713	3012
Genital Warts	285	222	507	1149	926	2075
Chancroid	0	0	0	0	0	0
Trichomoniasis	2	8	10	10	55	65
Candidiasis	291	421	712	1123	1544	2667
Bacterial Vaginosis	0	365	365	0	1368	1368
Other sexually transmitted disease	74	47	121	321	197	518
Non venereal <sup>8</sup>	683	456	1139	2845	1796	4641

<sup>\*</sup> Central STD clinic Colombo and peripheral STD clinics of National STD/AIDS Control Programme of Sri Lanka

- 1-Includes AIDS cases
- 2-Diagnosed within 2 years of infection and considered to be infectious.
- 3-Diagnosed after 2 years of infection and considered to be non-infectious
- 4-Includes both early and late cases
- 5-Includes presumptive Gonorrhoea
- 6-Includes both gonococcal and chlamydial conjunctivitis in neonatal period
- 7-Includes Lympho granuloma venerium, Granuloma inguinalae, Molluscum contagiosum, Scabis, Tinea, Hepatitis-Betc
- 8. Number of STD clinic attendees who were not having sexually transmitted diseses.

<sup>#</sup> Includes adjustments for revised diagnosis, reporting delays or any other amendments