## NATIONAL STD/AIDS CONTROL PROGRAMME

## QUARTERLY RETURN OF SEXUALLY TRANSMITTED DISEASE INCLUDING HIV/AIDS FOR EPIDEMIOLOGICAL BULLETIN OF THE EPIDEMIOLOGY UNIT OF MINISTRY OF HEALTH

New episodes of STD/HIV/AIDS reported or treated at STD clinics in Sri Lanka Year 2016 3rd Quarter

Disease	New cases or new disease episodes during the quarter			Total new cases or new episodes for calender year upto end of the quarter#		
	Male	Female	Total	Male	Female	Total
HIV Positives <sup>1</sup>	50	16	66	145	48	193
AIDS	13	3	16	36	9	45
Early Syphilis <sup>2</sup>	20	11	31	73	29	102
Late Syphilis <sup>3</sup>	143	72	215	363	213	576
Congenital Syphilis <sup>4</sup>	1	2	3	3	4	7
Gonorrhoea	45	11	56	172	51	223
Ophthalamia Neonatorum <sup>6</sup>	2	0	2	2	1	3
Non specific cervicitis/urethritis	150	404	554	435	1144	1579
Chlamydial infection	4	6	10	4	9	13
Genetal Herpes	331	426	757	1000	1295	2295
Genital Warts	329	242	571	864	704	1568
Chancroid	0	0	0	0	0	0
Trichomoniasis	0	9	9	8	47	55
Candidiasis	279	379	658	832	1123	1955
Bacterial Vaginosis	0	331	331	0	1003	1003
Other sexually transmitted disease	66	56	122	247	150	397
Non venereal <sup>8</sup>	772	423	1195	2162	1340	3502

<sup>\*</sup> Central STD clinic Colombo and peripheral STD clinics of National STD/AIDS Control Programme of Sri Lanka

# Includes adjustments for revised diagnosis , reporting delays or any other amendments

- 1-Includes AIDS cases
- 2-Diagnosed within 2years of infection and considered to be infectious.
- 3-Diagnosed after 2 years of infection and considered to be non-infectious
- 4-Includes both early and late cases
- 5-Includes presumptive Gonorrhoea
- 6-Includes both gonococcal and chlamydial conjunctivitis in neonatal period
- 7-Includes Lympho granuloma venerium, Granuloma inguinalae, Molluscum contagiosum, Scabis, Tinea, Hepatitis-B etc
- 8. Number of STD clinic attendees who were not having sexually transmitted diseses.