

## NATIONAL STD/AIDS CONTROL PROGRAMME

### QUARTERLY RETURN OF SEXUALLY TRANSMITTED DISEASE INCLUDING HIV/AIDS FOR EPIDEMIOLOGICAL BULLETIN OF THE EPIDEMIOLOGY UNIT OF MINISTRY OF HEALTH

<b>New episodes of STD/HIV/AIDS reported or treated at STD clinics in Sri Lanka Year 2017 1st Quarter</b>
---

Disease	New cases or new disease episodes during the quarter			Total new cases or new episodes for calender year upto end of the quarter#		
	Male	Female	Total	Male	Female	Total
HIV Positives <sup>1</sup>	56	17	73	56	17	73
AIDS	6	2	8	6	2	8
Early Syphilis <sup>2</sup>	18	2	20	18	2	20
Late Syphilis <sup>3</sup>	130	59	189	130	59	189
Congenital Syphilis <sup>4</sup>	1	1	2	1	1	2
Gonorrhoea	69	18	87	69	18	87
Ophthalmia Neonatorum <sup>6</sup>	0	1	1	0	1	1
Non specific cervicitis/urethritis	149	485	634	149	485	634
Chlamydial infection	1	0	1	1	0	1
Genetal Herpes	293	443	736	293	443	736
Genital Warts	285	253	538	285	253	538
Pelvic inflammatory disease		17	17		17	17
Trichomoniasis	1	13	14	1	13	14
Candidiasis	253	414	667	253	414	667
Bacterial Vaginosis		351	351		351	351
Other sexually transmitted disease <sup>7</sup>	72	35	107	72	35	107
Non venereal <sup>8</sup>	662	641	1303	662	641	1303

\* Central STD clinic Colombo and peripheral STD clinics of National STD/AIDS Control Programme of Sri Lanka

# Includes adjustments for revised diagnosis , reporting delays or any other amendments

1-Includes AIDS cases

2-Diagnosed within 2years of infection and considered to be infectious.

3-Diagnosed after 2 years of infection and considered to be non-infectious

4-Includes both early and late cases

5-Includes presumptive Gonorrhoea

6-Includes both gonococcal and chlamydial conjunctivitis in neonatal period

7-Includes Lympho granuloma venerium, Granuloma inguinale, Molluscum contagiosum, Scabis, Tinea, Hepatitis- B etc

8-.Number of STD clinic attendees who were not having sexually transmitted diseases.