## NATIONAL STD/AIDS CONTROL PROGRAMME

## QUARTERLY RETURN OF SEXUALLY TRANSMITTED DISEASE INCLUDING HIV/AIDS FOR EPIDEMIOLOGICAL BULLETIN OF THE EPIDEMIOLOGY UNIT OF MINISTRY OF HEALTH

## New episodes of STD/HIV/AIDS reported or treated at STD clinics in Sri Lanka Year 2018 3rd Quarter

	New cases or new disease			Total new cases or new episodes		
Disease	episodes during the quarter			for the calendar year up to end		
				of the quarter#		
	Male	Female	Total	Male	Female	Total
HIV Positives1	79	20	99	223	52	275
AIDS	15	6	21	41	9	50
Early Syphilis2	17	5	22	36	8	44
Late Syphilis3	114	67	181	379	188	567
Congenital Syphilis4	1	0	1	2	1	3
Gonorrhoea5	55	29	84	145	59	204
Ophthalmia Neonatorum6	0	0	0	1	0	1
Non specific cervicitis/urethritis	235	524	759	647	1431	2078
Chlamydial Infection	1	0	1	7	2	9
Genital Herpes	335	484	819	963	1369	2332
Genital Warts	353	308	661	1003	834	1837
Pelvic inflammatory disease	0	32	32	0	79	79
Trichomoniasis	1	16	17	7	35	42
Candidasis	337	488	825	955	1431	2386
Bacterial Vaginosis	0	418	418	0	1118	1118
Other sexually transmitted diseas	145	88	233	376	224	600
Non-venereal8	954	526	1480	2722	1568	4290

<sup>\*</sup> Central STD clinic Colombo and peripheral STD clinics of National STD/AIDS Control Programme of Sri Lanka

- 2-Diagnosed within 2 years of infection and considered to be infectious.
- 3-Diagnosed after 2 years of infection and considered to be non-infectious
- 4-Includes both early and late cases
- 5-Includes presumptive Gonorrhoea
- 6-Includes both gonococcal and chlamydial conjunctivitis in neonatal period
- $\hbox{\it 7-Includes Lympho granuloma venerium, Granuloma inguinalae,} Mollus cum contagios um, Scabis, Tinea, Hepatitis-Betc.$
- 8-Number of STD clinic attendees who were not having sexually transmitted diseases

<sup>#</sup> Includes adjustments for revised diagnosis, reporting delays or any other amendments

<sup>1-</sup>Includes AIDS cases